




Advanced Research in  
Microbial Metabolites & Technology  
(ARMMT)

journal homepage: <http://armmt.irost.ir>



## Heterologous expression and characterization of a recombinant thermostable amylopullulanases *Coh4159* from *Cohnella* sp. A01

Parvin Valiulahi<sup>1</sup>, Saeed Aminzadeh<sup>1\*</sup>, Mehdi Shamsara<sup>2</sup>, Naser Farrokhi<sup>3</sup>, Jahan Alikhajeh<sup>4</sup>

<sup>1</sup>Bioprocess Engineering Research Group, Institute of Industrial and Environmental Biotechnology, National Institute for Genetic Engineering and Biotechnology, Tehran, Iran

<sup>2</sup>Animal Biotechnology Department, Institute of Agricultural Biotechnology, National Institute for Genetic Engineering and Biotechnology, Tehran, Iran

<sup>3</sup>Department of Plant Biology & Biotechnology, Faculty of Life Sciences & Biotechnology, Shahid Beheshti University G.C., Evin, Tehran, Iran

<sup>4</sup>Departments of Physiology and Cellular Biophysics Columbia University Medical Center, USA

### Article Info

Received: 22 January 2018

Received in revised form:  
24 June 2018

Accepted: 09 July 2018

### Keywords:

amylopullulanase  
pullulan  
recombinant protein  
zymogram

### Abstract

Starch debranching enzymes that merely hydrolyze  $\alpha$ -(1 $\rightarrow$ 6) glycosidic linkages are classified into isoamylases (EC 3.2.1.68) and pullulanases (EC 3.2.1.41). An exception to this definition is amylopullulanase, a type of pullulanase that is capable of cleaving both  $\alpha$ -(1 $\rightarrow$ 4) and  $\alpha$ -(1 $\rightarrow$ 6) linkages. Amylopullulanases are in demand in liquid sugar industries to generate glucose and some other starch derivatives. Pullulanases can be used in conjunction with amylases to improve sugar availability during sugar syrup production. Here, a thermophilic *Cohnella* sp. A01 amylopullulanase (EC 3.2.1.41) gene, namely *Coh4159*, was PCR amplified and cloned in pET-26b(+) and transformed into BL21(DE3). Recombinant *Coh4159* was heterologously expressed in the presence of 0.5 mM IPTG and purified via affinity chromatography, and further characterized. Enzyme activity was demonstrated via zymogram analysis in the presence of pullulan. The enzyme had a hydrolytic effect on pullulan with  $V_{\max} = 2.85 \mu\text{mol}\cdot\text{min}^{-1}$  and  $K_m = 0.5 \text{ mM}$ . Temperature optima and pH were 60 °C and 6.0, in which the enzyme kept its activity at wide pH (4-9) and temperature (30-70 °C) ranges. The recombinant enzyme kept 50% of its activity for 60 min, 100 min and 120 min when incubated at 80, 70 and 60 °C, respectively. Amongst metal ions tested,  $\text{Mn}^{2+}$  and  $\text{Ca}^{2+}$  have improved the enzyme activity both at 5 and 10 mM. The results promise the capability of producing a commercial industrial enzyme well-suited to liquid sugar syrup industry specifications.

## 1. Introduction

The sugar industry is a thriving business that mainly relies on agriculture with a focus on sugarcane and sugar beet. These crop sources barely satisfy daily human consumption around the globe, a reason that justifies the

increasing price of sugars in the last few years. The growing demand on sugar in other food and drug industries provides enough incentive to seek other natural sources. Starch, as a byproduct of industrial crops, such as cassava, potato and corn, has proven to be a valuable source to produce glucose. Acid hydrolysis of starch has been used for its conversion to

\*Corresponding author. Tel: +982144787412

E-mail address: aminzade@nigeb.ac.ir

DOI: 10.22104/ARMMT.2018.2738.1014

smaller reduced sugars and glucose. However, use of chemicals in such refineries makes the final product expensive due to high energy demand, the factory sewage also contaminates the environment. In contrast, enzymatic bioconversion of starch can efficiently turn the biopolymer to sugar syrups containing glucose, dextrans and other oligosaccharides (Janeček *et al.*, 2014) with minimal environmental effects and cost.

Starch is a polymer of  $\alpha$ -(1 $\rightarrow$ 4) glucosyl residues, found both in amylose and amylopectin, that is substituted with chains of  $\alpha$ -(1 $\rightarrow$ 6) glucosyl units in amylopectin alone. Starch debranching enzymes that merely hydrolyze  $\alpha$ -(1 $\rightarrow$ 6) glycosidic linkages are classified into isoamylases (EC 3.2.1.68) and pullulanases (EC 3.2.1.41). An exception to this definition is amylopullulanase, a type of pullulanase that is capable of cleaving both  $\alpha$ -(1 $\rightarrow$ 4) and  $\alpha$ -(1 $\rightarrow$ 6) linkages (Wang *et al.*, 2012). The difference between these two lies in the fact that the pullulanases prefer pullulan as the substrate (Martin *et al.*, 1995; van der Veen *et al.*, 2000). Pullulan,  $\alpha$ -(1 $\rightarrow$ 4),  $\alpha$ -(1 $\rightarrow$ 6) glucan is a linear homopolysaccharide of 3 glucosyl residues, maltotriose, with  $\alpha$ -(1 $\rightarrow$ 6) linkages that are connected to each other by an  $\alpha$ -(1 $\rightarrow$ 4) bond. This water-soluble tasteless polymer has many food, drug and industrial applications (Leathers, 2003).

Pullulanases can be used in conjunction with amylases to improve sugar availability during sugar syrup production (Hii *et al.*, 2012). Likewise, in the baking industry bread hardening and taste change can be slowed down by using amylopullulanases instead of polyoxyethylene monostearate to produce short chain water soluble dextrans (Lord, 1950). These molecules can change the nature of starch gelatinization and therefore slow down bread staling (Prakash *et al.*, 2012). For instance, supplementation of whole wheat dough with an amylopullulanase from *Geobacillus* improved the loaf volume, shelf-life, and bread texture. Additionally, the

enzyme was capable of desizing cotton fabrics (Nisha and Satyanarayana, 2014). Pullulanases are also being used in cellodextrin formation which has application in pharmaceutical industries to improve solubility of water-insoluble drugs, to add to skin protectants against UV light, and in food industries to cover wrong flavor (Hassan *et al.*, 2006; Prakash *et al.*, 2012). Furthermore, pullulanases have a stake in the biofuel industry, allowing the release of more glucose from starch which can be harnessed during fermentation (KiliApar and Özbek, 2004; Prakash *et al.*, 2012; Nahampun *et al.*, 2013). For example, the ectopic expression of *Thermoanaerobacter* amylopullulanase in maize seeds under the control of a seed specific promoter improved the starch conversion efficiency by 90.5%, enhancing bioethanol production (Nahampun *et al.*, 2013). Last but not least, amylopullulanases have found their ways into agricultural biotechnology through transgenesis of cereals to accelerate bioprocessing of starch and producing protein-enriched flour (Chiang *et al.*, 2005). Despite better bioavailability of amino acids in transgenic rice, the plants grown in farm demonstrated higher susceptibility to brown spot caused by *Bipolaris oryzae* (Ting *et al.*, 2008).

Since more than 90% of industrial enzymes are mainly obtained from microorganisms, (Cho *et al.*, 2007), looking for new enzyme sources with extraordinary features are always a blessing for the industry. Thermophilic bacteria bear such enzymes that are highly tolerant to extreme temperatures (Narang *et al.*, 2001), something that is inevitable in technologies currently in use (Mrudula *et al.*, 2011). We have previously reported the isolation of *Cohnella* A01, a Gram positive thermophilic bacterium, from a shrimp pond (Aliabadi *et al.*, 2016), and heterologous expression of its thermostable lipase (Pooreydy Golaki *et al.*, 2015a), structural characterization of lipase (PooreydyGolaki *et al.*, 2015b), cloning of  $\beta$ -glucanase (Rezaie *et al.*, 2018) and two other amylopullulanases of this bacterium (Zebardast *et al.*, 2017). Here an amylopullulanase sequence

(GenBank accession: KX013443), namely *Coh4159*, from its fully sequenced genome (data not shown) was found through in silico studies and used for heterologous expression and purification followed by enzymatic assays.

## 2. Materials and methods

### 2.1. Materials

All enzymes, pET-26b(+), IPTG and X-Gal were purchased from Fermentas (Burlington, Canada). DNA extraction kit was obtained from BioNEER (Anaheim, USA). High Pure PCR Product Purification and High Pure Plasmid Purification kits were obtained from Roche (Basel, Switzerland). *E. coli* host strains, DH5 $\alpha$  and BL21 (DE3), were purchased from Invitrogen (Carlsbad, CA, USA). Ni-NTA resin was obtained from Qiagen (Netherlands). Other materials were from Merck (Darmstadt, Germany). *Cohnella* sp. A01 was obtained from sampled water from a shrimp pond at Choebdeh (Abadan, Iran; Aliabadi *et al.*, 2016).

### 2.2. Polymerase Chain Reaction

Single clone of *Cohnella* sp. A01 was cultured in nutrient broth for 72 h at 60 °C. The culture was centrifuged at 7000  $\times$ g for 10 min. DNA was isolated from precipitated bacterial cells using the DNA extraction kit as instructed by the supplier. DNA (70 ng/ $\mu$ l) was used for PCR amplification of amylopullulanase in a 25  $\mu$ l reaction containing 10 pmoles of each primers [(Forward:5'-CTTCCATGGTGAA CAAGCAACTGGGCAAATCCTCGC-3', underline is *Nco*I restriction site and Reverse:5'-CTTCTCGAGTGGACGCTCCAGCAGCCG GAC-3', underlined is *Xho*I restriction site)], 0.5  $\mu$ l of 100 U Pfu DNA polymerase, 2.5  $\mu$ l of 10 $\times$  PCR buffer, 0.5  $\mu$ l of 10 mM dNTPs and 1.5  $\mu$ l DMSO. The PCR was started at 94 °C for 5 min then followed by 35 cycles of

94 °C: 30 s; 61.5 °C: 30 s; 72 °C: 2 min with a final extension at 72 °C for 10 min. The PCR product (2439 bp) was separated on 1% agarose gel and isolated with a High Pure PCR Product Purification kit according to the manufacturer. The PCR product was restricted with 10 U/ $\mu$ l of *Nco*I and *Xho*I, separated on 1% agarose gel and purified as mentioned above and named *Coh4159*.

### 2.3. Cloning and expression of *Coh4159*

pET-26b(+) plasmid vector supplied in the bacterial cell culture was isolated using a High Pure Plasmid Purification kit according to the manufacturer. The plasmid was restricted with *Nco*I and *Xho*I as stated earlier at 37 °C for 2 h, separated on 1% agarose gel and purified using a High Pure PCR Product Purification kit according to the manufacturer. *Coh4159* (25 ng/ $\mu$ l) was ligated with pET-26b(+) (37 ng/ $\mu$ l) using T4-DNA ligase (5 U/ $\mu$ l) in the presence of 1  $\mu$ l 10 $\times$  T4-DNA ligase buffer at 22 °C for 2 h and 1 h at 25 °C. The construct was used to transform competent BL21 cells via the heat-shock method (Green and Sambrook, 2012). The cloned plasmid, pET-Amylo, was isolated as mentioned above and sequenced using M13 primers for sequence confirmation.

BL21 cells containing pET-Amylo were cultured in 5 ml LB containing 25  $\mu$ g/ml kanamycin for 16 h at 37 °C. The culture was used to seed a 50 ml LB until OD<sub>600</sub> = 0.6. IPTG (0.5 mM) was used to induce the T7 promoter at 25 °C on a rotary shaker at 150 rpm for 6 h. The bacterial cells expressing pET-Amylo were precipitated at 6000  $\times$ g for 10 min at 4 °C. The cells were resuspended in lysis buffer (50 mM NaH<sub>2</sub>PO<sub>4</sub>, 500 mM NaCl, 10 mM imidazole, 0.05% v/v Tween20) and incubated for 45 min at 4 °C. The cells were broken open using a sonifier at 70% strength and five 0.5 pulses on ice. The mixture was centrifuged at 6000  $\times$ g for 10 min at 4 °C. The supernatant was used for recombinant protein purification via affinity

chromatography using Ni-NTA resin according to the manufacturer. The isolated recombinant protein was dialyzed for 24 h against 20 mM  $\text{NaH}_2\text{PO}_4$  (pH = 7.5) at 4 °C with buffer change every 8 h.

#### 2.4. Activity assays for recombinant amylopullulanases

Amylopullulanase activity was determined according to the 3,5-dinitrosalicylic acid (DNS) method (Bernfeld, 1955). The substrate was 1% pullulan in mixed buffer (50 mM of each sodium phosphate, sodium acetate and glycine, pH = 5.5). To assay the protein activity, enzyme (50  $\mu\text{l}$ ) was added to the mixture of 50  $\mu\text{l}$  1% pullulan and incubated for 20 min at 60 °C. DNS reagent (100  $\mu\text{l}$  of 1% w/v DNS, 30% w/v sodium potassium tartarate, 1.6% w/v sodium hydroxide) was added to stop the progression of reaction. The reaction was incubated at 100 °C for 10 min and after cooling down to 22 °C, centrifuged at 6000  $\times g$  for 10 min, and the production of reduced sugars was read at 530 nm. By definition, 1 unit of amylopullulanase can release 1  $\mu\text{mole}$  of reduced sugar in 1 min and in our case at 60 °C (pH = 6.0), when maltose is being used as the standard.

#### 2.5. Definition of Biochemical and kinetic parameters

A pH profile was established with 1% pullulan in 50 mM mixed buffer at pHs 3-12 and an enzyme assay was performed as mentioned above. For pH stability, enzymes were incubated for 3 h at pH 3, 7 and 11 in 50 mM mixed buffer. The substrate prepared at optimum pH was added to the enzyme and the activity was measured as stated earlier. Furthermore, relative enzyme activities were determined in temperatures ranging from 20-90 °C. For this, 200  $\mu\text{l}$  enzyme with 200  $\mu\text{l}$  substrate were incubated separately at the temperatures for 60 min and enzyme activity was assayed. For

temperature stability measurement, 600  $\mu\text{l}$  of enzyme was incubated at 60, 70 and 80 °C for 3 h. Every 10 min, 20  $\mu\text{l}$  of amylopullulanase incubated in different temperatures, was added to 20  $\mu\text{l}$  substrate and the enzyme activity was assayed.

#### 2.6. Effects of chemical compounds and metal ions on Coh4159 activity

Metal cations, 5 mM of each  $\text{MgCl}_2$ ,  $\text{CaCl}_2$ ,  $\text{KCl}$ ,  $\text{FeCl}_2$ ,  $\text{MnCl}_2$ ,  $\text{ZnCl}_2$ ,  $\text{CuCl}_2$ , and  $\text{NaCl}$ , were analyzed to check the effect on the enzyme activity.  $V_{\text{max}}$  and  $K_m$  were established for both recombinant enzymes with Lineweaver-Burk (Cook and Cleland, 2007) using pullulan as substrate. Every experiment had three replicates and the data were plotted in Prism5 (Motulsky, 2007).

#### 2.7. Zymogram analysis

Native-PAGE (10%) including pullulan was used to separate the isolated recombinant enzyme. Triton X100 (2.5%) was used to wash the gel for 1 h with 50 mM sodium acetate (pH = 6.0) at 4 °C. The gel was incubated at 60 °C for 1 h then Lugol's reagent was added to the gel and enzyme activity was photographed.

---


### 3. Results and discussion

#### 3.1. Expression and purification of the recombinant amylopullulanase

The *Coh4159* gene was PCR amplified, restricted with *NcoI* and *XhoI* and cloned in pET-26b(+) and transformed into BL21(DE3). Recombinant Coh4159 was heterologously expressed in the presence of 0.5 mM IPTG and purified via affinity chromatography (Figure 1, lane 3). The enzyme relative molecular weight was 82, kDa which can be considered as a relatively small to medium size



amylopullulanases (Table 1). Enzyme activity was demonstrated via zymogram analysis in the presence of pullulan (Figure 1, lane 5).



**Figure 1.** Heterologous expression of recombinant Coh4159 in *E. coli* and zymogram analysis. Lane 1: un-induced bacterial cell culture, Lane 2: induced bacterial cell culture with prominent band around 82 kDa. Lane 3: purified Coh4159 via affinity chromatography using Ni-NTA column, Lane 4: protein ladder, Lane 5: zymogram analysis is illustrative of enzyme activity in hydrolyzing pullulan.

### 3.2. Coh4159 kinetic and biochemical parameters

Enzymes kinetic parameters in the presence of pullulan were calculated with Prism5. Coh4159 had  $V_{max} = 2.85 \mu\text{mol}\cdot\text{min}^{-1}$  and  $K_m = 0.5 \text{ mM}$  (Figure 2a). Optimum pH for Coh4159 was 6.0 (Figure 2b), and the enzyme kept 50% of its activity for 2 h at pH = 7.0 and for 50 min at pH 3, and 11 (Figure 2c). Thus, the pH optima was similar to other previously reported amylopullulanases, except for Zebardast *et al.* (2017) where the greatest pH activity was obtained at basic pH = 8.0. The highest activity for Coh4159 was seen at 60 °C (Figure 2d). The recombinant enzyme kept 50% of its activity for 60 min, 100 min and 120 min incubated at 80, 70 and 60 °C, respectively (Figure 2e). Therefore, the enzyme had great temperature stability when compared to homologous enzymes from other bacterial sources. Although, some other amylopullulanases with greater tolerance to higher temperature have been previously reported (Table 1).

**Table 1.** Some properties of characterized amylopullulanases from thermophilic bacteria


Bacterial species	MW <sup>†</sup> (kDa)	T. O <sup>‡</sup> (°C)	pH <sup>#</sup>	Ca <sup>2+Δ</sup>	Reference
<i>Bacillus flavocaldarius</i> KP1228	55	75-80	6.3	-	Suzuki <i>et al.</i> , 1991
<i>Pyrococcus furiosus</i>	89	105	5.5	d	Dang <i>et al.</i> , 1997
<i>Thermoanaerobacterium thermosaccharolyticum</i>	150	60	5-5.5	-	Ganghofner <i>et al.</i> , 1998
<i>Thermoanaerobacterium ethanolicus</i> 39E	109	80	6.0	d	Lin and Leu, 2002
<i>Staphylothermus marinus</i>	73.5	105	5	d	Li <i>et al.</i> , 2013
<i>Geobacillus thermoleovorans</i> (gt-apu)	182	60	7.0	-	Nisha and Satyanarayana, 2013
<i>Geobacillus thermoleovorans</i> (apu105)	105	80	7.0	i	Nisha and Satyanarayana, 2014
Coh00831	70	60	6.0	d	Zebardast <i>et al.</i> , 2017
Coh01133	127	70	8.0	d	Zebardast <i>et al.</i> , 2017
Coh4159	82	60	6.0	d	This study

<sup>†</sup> Molecular weight

<sup>‡</sup> Temperature optimum

<sup>#</sup> pH optimum


<sup>Δ</sup> Calcium dependency, d: dependent, i: independent



**Figure 2.** Kinetic parameters of Coh4159. a) Michaelis-Menton graph, b) pH profile, c) pH stability, d) temperature profile, and e) temperature stability.

### 3.3. Metal ions and chemical compounds effects

Amongst metal ions tested,  $Mn^{2+}$  and  $Ca^{2+}$  have improved the enzyme activity both at 5 and 10 mM. So far the only reported  $Ca^{2+}$  independent amylopullulanase was from *Geobacillus thermoleovorans* (Nisha and Satyanarayana, 2014), while similar to this study the others were dependent (Table 1).  $Mg^{2+}$  increased the enzyme activity at 5 mM, while 10 mM  $Mg^{2+}$  was no different than the control. Other cations significantly decreased the enzyme activity (Figure 3).



**Figure 3.** The effect of 5 and 10 mM cations on amylopullulanase activity. The thin bars are 10 mM of each metal cations.

## 4. Conclusion

Due to high amylopullulanase activity and thermostability, we found the recombinant enzyme suitable for use in industries such as baking, liquid sugar and bioethanol. In the future, we plan to transform the plasmid in an industrial strain and further optimize the culture condition for greater production of the recombinant enzyme.

## Conflict of interest

The authors declare that there is no conflict of interest.

## Acknowledgements

We would like to thank National Institute of Genetic Engineering and Biotechnology (NIGEB) for providing the research facility.

## Ethical approval

This manuscript does not contain any studies with human participants or animals performed by any of the authors.

## Open access

This article distributed under the terms of the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## References

- [1] Aliabadi N, Aminzadeh S, Karkhane A A, Haghbeen K (2016) Thermostable chitinase from *Cohnella* sp. A01:isolation and product optimization. *Brazilian Journal of Microbiology*. 47: 931-940. doi: 10.1016/j.bjm.2016.07.009
- [2] Bernfeld P (1995) Amylase  $\alpha$  and  $\beta$ In. *Methods in Enzymology* (ed. Colowick S, Kaplan, N), 1: 149-158. [http://dx.doi.org/10.1016/0076-6879\(55\)01021-5](http://dx.doi.org/10.1016/0076-6879(55)01021-5)

- [3] Chiang C-M, Y F-S, Huang L-F, Tseng T-H, Chung M-C, Wang C-S, Lur H-S, Shaw J-F, Yu S-M (2005) Expression of a bi-functional and thermostable amylopullulanase in transgenic rice seeds leads to autohydrolysis and altered composition of starch. *Molecular Breeding*. 15: 125. doi: 10.1007/s11032-004-3919-7
- [4] Cook PF, Cleland WW (2007) *Enzyme kinetics and mechanism*. Garland Science New York. doi: 10.1110/ps.073353008
- [5] Green MR, Sambrook J (2001). Cloning and transformation with plasmid vectors. In. *Molecular cloning: A laboratory manual*. Fourth edition. CSHL, New York, USA. 157-258.
- [6] Dong G, Vieille C, Zeikus JG (1997) Cloning, sequencing, and expression of the gene encoding amylopullulanase from *Pyrococcus furiosus* and biochemical characterization of the recombinant enzyme. *Applied and Environmental Microbiology*. 63: 3577-3584.
- [7] Ganghofner D, Kellermann J, Staudenbauer WL, Bronnenmeier K (1998) Purification and properties of an amylopullulanase, a glucoamylase, and an  $\alpha$ -glucosidase in the amylolytic enzyme system of *Thermoanaerobacterium thermosaccharolyticum*. *Biosci. Biotechnol. Biochem*. 62: 302-308.
- [8] Hasan F, Shah AA, Hameed A (2006) Industrial applications of microbial lipases. *Enzyme and Microbial Technology*, 39: 235-251. <https://doi.org/10.1016/j.enzmictec.2005.10.016>.
- [9] Hii SL, Tan JS, Ling TC, Ariff AB (2012) Pullulanase: role in starch hydrolysis and potential industrial applications. *Enzyme Res*. doi: 10.1155/2012/921362. doi: 10.1155/2012/921362.
- [10] Janeček Š, Svensson B, MacGregor EA. (2014).  $\alpha$ -Amylase: an enzyme specificity found in various families of glycoside hydrolases. *Cellular and Molecular Life Sciences*, 71(7): 1149-1170. doi: 10.1007/s00018-013-1388-z.
- [11] Kılıç Apar D, Özbek B (2004)  $\alpha$ -Amylase inactivation by temperature during starch hydrolysis. *Process Biochemistry*. 39(9): 1137-1144.
- [12] Leathers TD (2003) Bioconversions of maize residues to value-added coproducts using yeast-like fungi. *FEMS Yeast Research*. 3(2): 133-140.
- [13] Li X, Li D, Park KH (2013) An extremely thermostable amylopullulanase from *Staphylothermus marinus* displays both pullulan and cyclodextrin-degrading activities. *Appl Microbiol Biotechnol*. 97: 5359-5369. doi: 10.1007/s00253-012-4397-1.
- [14] Lin FP, Leu KL (2002) Cloning, expression, and characterization of thermostable region of amylopullulanase gene from *Thermoanaerobacter ethanolicus* 39E. *Appl. Biochem. Biotechnol*. 97: 33-44.
- [15] Lord DD (1950) The action of polyoxyethylene monostearate upon starch with reference to its softening action in bread. *J. Colloid Sci*. 5: 360.
- [16] Martin C, Smith, AM (1995) Starch biosynthesis. *The Plant Cell*. 7(7): 971.
- [17] Motulsky H (2007) *Prism 5 Statistics Guide*. GraphPad Software.
- [18] Mrudula S, Reddy, G Seenayya G (2011) Effect of process parameters on production of thermostable amylopullulanase by *Clostridium thermosulfurogenes* SVM 17 under solid state fermentation. *Malaysian Journal of Microbiology* 7(1): 19-25.
- [19] Nahampun HN, Lee CJ, Jane JL, Wang K (2013) Ectopic expression of bacterium amylopullulanase enhances bioethanol production from maize grain. *Plant Cell Rep*. 32: 1393-1405. doi: 10.1007/s00299-013-1453-y.
- [20] Narang S, Satyanarayana T (2001) Thermostable  $\alpha$ -amylase production by an extreme thermophile *Bacillus thermoleovorans*. *Letters in Applied Microbiology*, 32(1): 31-35.
- [21] Nisha M, Satyanarayana T (2013) Characterization of recombinant amylopullulanase (gt-apu) and truncated amylopullulanase (gt-apuT) of the extreme thermophile *Geobacillus thermoleovorans* NP33 and their action in starch. *Appl Microbiol Biotechnol*. 97: 6279-6292. doi: 10.1007/s00253-012-4538-6.
- [22] Nisha M, Satyanarayana T (2014) Characterization and multiple applications of a highly thermostable and  $\text{Ca}^{2+}$ -independent amylopullulanase of the extreme thermophile *Geobacillus thermoleovorans*. *Appl Biochem Biotechnol*. 174: 2594-2615.
- [23] Pooreydy Golaki B, Aminzadeh S, Karkhane A A, Yakhchali B, Farrokh P, Khaleghinejad S H, Akhavian Tehrani A, Mehrpooyan S. (2015a) Cloning, expression, purification, and characterization of lipase 3646 from thermophilic indigenous *Cohnella* sp. A01. *Protein Expression and Purification* 109: 120-126. doi: 10.1016/j.pep.2014.10.002

- [24] Pooreydy Golaki B, Aminzadeh S, Karkhane A A, Yakhchali B, Farrokh P, RastgarJazii F, Nadimifar M. (2015b) The Effect of pH on Globular State of Lipase-3646; an Appropriate Model for Molten Globule Investigations. *Protein J.* 34:267–274. doi: 10.1007/s10930-015-9622-1.
- [25] Prakash N, Gupta S, Ansari M, Ahmad Khan Z, Suneetha V (2012) Production of economically important products by the use of pullulanase enzyme. *Int J Sci Innovat Discover*, vol.2. pp.266-273.
- [26] Rezaie M, Aminzadeh S, Heidari F, Mashhadi Akbar Boojari M, Karkhane A A (2018) Biochemical Characterization of Recombinant Thermostable *Cohnella* sp. A01  $\beta$ -Glucanase. *Iranian Biomedical Journal*, ahead of publication.
- [27] Suzuki Y, Hatagaki K, Oda H (1991) A hyperthermostable pullulanase produced by an extreme thermophile, *Bacillus flavocalfarius* KP 1228, and evidence for the proline theory of increasing protein thermostability. *Appl Microbiol Biotechnol.* 34: 707-714.
- [28] Ting M-Y, Shih H-D Lin C-Y (2008) Increased susceptibility of rice following insertion of amylopullulanase gene, to brown spot caused by *Bipolaris oryzae*. *J. Phytopathol.* 156: 530-533.
- [29] van der Veen BA, Uitdehaag J, Dijkstra BW, Dijkhuizen L (2000) Engineering of cyclodextrin glycosyltransferase reaction and product specificity. *Biochimica et Biophysica Acta (BBA)-Protein Structure and Molecular Enzymology*, 1543(2): 336-360.
- [30] Wang L-Y, Chen S-F, Wang, L, Zhou Y-G, Liu H-C (2012) *Cohnella plantaginis* sp. nov., a novel nitrogen-fixing species isolated from plantain rhizosphere soil. *Antonie van Leeuwenhoek*, 102(1): 83-89. doi: 10.1007/s10482-012-9715-1
- [31] Zebardast Roodi F, Aminzadeh S, Farrokhi N, Karkhane A A, Haghbeen K (2017) *Cohnella* amylopullulanases: Biochemical characterization of two recombinant thermophilic enzymes. *PLOS ONE* 12(4): 1-13. <https://doi.org/10.1371/journal.pone.0175013>